

Static response of Fermi liquids with tensor interactions

E. Olsson

*Department of Astronomy and Space Physics, Uppsala University, Box 515, 75120 Uppsala, Sweden
and NORDITA, Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark*

P. Haensel

Nicolaus Copernicus Astronomical Center, Bartycka 18, 00-716 Warsaw, Poland

C. J. Pethick

NORDITA, Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark

(Dated: February 8, 2008)

We use Landau's theory of a normal Fermi liquid to derive expressions for the static response of a system with a general tensor interaction that conserves the total spin and the total angular momentum of the quasiparticle-quasihole pair. The magnetic susceptibility is calculated in detail, with the inclusion of the center of mass tensor and cross vector terms in addition to the exchange tensor one. We also introduce a new parametrization of the tensor Landau parameters which significantly reduces the importance of high angular harmonic contributions. For nuclear matter and neutron matter we find that the two most important effects of the tensor interaction are to give a contribution from multipair states and to renormalize the magnetic moments. Response to a weak probe may be calculated using similar methods, replacing the magnetic moments with the matrix elements of the weak charges.

I. INTRODUCTION

For systems with central interactions, Landau's theory of normal Fermi liquids provides an economical way of characterizing many low-temperature properties. The theory was applied to atomic nuclei by Migdal and collaborators, and in that work it was generally assumed that the non-central contributions to the effective nucleon-nucleon interaction were small [1]. The generalization of Landau theory to include effects of the tensor force was made by Dąbrowski and Haensel [2, 3, 4]. Subsequently, estimates of tensor contributions to the effective interaction were made in Refs. [5, 6, 7, 8].

The stimulus for the present work arose in the context of astrophysics. In the physics of collapse and the subsequent evolution of a neutron star, the properties of neutrinos in dense matter are a key ingredient [9]. As demonstrated in Refs. [10, 11, 12, 13, 14, 15, 16, 17], neutrino scattering and absorption rates are sensitive to nucleon-nucleon interactions, especially their spin-dependent parts. Direct calculation of effective interactions is difficult, but for systems with only central interactions, information about the effective interaction at long wavelengths may be obtained directly from a knowledge of static properties of the matter. For example, the spin susceptibility of a single-component Fermi liquid with two spin states is given by

$$\chi = \frac{\mu_0^2 N(0)}{1 + G_0}, \quad (1)$$

where μ_0 is the magnetic moment of a particle in free space, $N(0) = m^* p_F / \pi^2 \hbar^3$ is the density of states per unit volume at the Fermi surface, m^* is the quasiparticle effective mass, p_F is the Fermi momentum and G_0 is the Landau parameter describing the isotropic part

of the spin dependent contribution to the quasiparticle interaction. Since calculations of the magnetic susceptibility of neutron matter exist [18], it is relevant to ask to what extent it is possible to deduce properties of the effective interaction from such data. A related question is the extent to which tensor contributions to quasiparticle energies and interactions alter neutrino scattering rates, which were previously calculated neglecting tensor effects in Ref. [11].

As shown in earlier work [19], the tensor force influences the static response of a system in a number of different ways. One is that the magnetic moment of a quasiparticle is different from its value for a particle in vacuum. In particular, the magnetic moment is not a scalar, as it is for systems with central forces only. As long ago as 1951, Miyazawa [20] calculated explicitly how the magnetic moment of a nucleon would be modified by the tensor interaction due to one pion exchange and more recent discussions may be found in e.g. the review [21] and Ref. [22]. A second effect is that the quasiparticle interaction contains explicit tensor contributions. Following Refs. [2, 3, 4], it has generally been assumed that these have an exchange-tensor structure similar to that of the one-pion exchange interaction. However, Schwenk and Friman [7] have pointed out recently that the one-pion exchange interaction, when acting in second order, can give rise to contributions to the effective interaction which have a different structure. In their paper they evaluated these induced interaction contributions to the quasiparticle interaction and found that the exchange-tensor term in the quasiparticle interaction is much reduced, and that terms of a different structure can be of a comparable magnitude to the exchange-tensor ones. A third effect of the tensor interaction is that there are multipair contributions to the magnetic susceptibility.

In this paper we begin by deriving a general expression for the magnetic susceptibility in Sec. III, taking into account tensor contributions to the magnetic moment, and tensor contributions to the effective interaction which are completely general for an interaction which conserves the total angular momentum and the total spin of the quasiparticle-quasihole pair. This represents a generalization of the earlier calculation of the magnetic susceptibility by Haensel and Jerzak [23], who took into account the effect of the exchange-tensor contribution to the quasiparticle interaction. Another issue that we address is how to parametrize tensor contributions to the quasiparticle interaction (Sec. IV). The scheme usually employed in the past suffers from the disadvantage that it is generally necessary to take into account high angular harmonic contributions to the interaction. We present an alternative parametrization for which higher harmonic terms play little role. In Sec. V we evaluate the matrix elements needed for the calculation of the susceptibility and estimate the magnitudes of the different contributions. In Sec. VI we extend the result to symmetric nuclear matter and to responses with other spin and isospin properties, for example that to weak interactions. Sec. VII contains concluding remarks.

II. BASIC FORMALISM

The reason for Landau's theory of normal Fermi liquids being particularly simple for systems with interparticle interactions that are central is that the energies of long-wavelength, low-lying states differing little from the ground state may be described solely in terms of the quasiparticle distribution. The importance of conservation laws in determining the low-frequency, long-wavelength behaviour of Fermi systems has been stressed by Leggett [24] and a discussion in terms of Landau theory is given in Ref. [25]. Expressed in other terms, the only excitations of importance are ones with a single added quasiparticle and a single quasihole. When non-central interactions are present, the energy must include contributions from states with more than one quasiparticle-quasihole pair. To facilitate satisfying the conservation law for particle number, it is convenient to work with the thermodynamic potential $\langle \hat{H} - \tilde{\mu} \hat{N} \rangle$ which, for brevity, we shall refer to simply as the energy. Here $\hat{H}$ is the Hamiltonian operator, $\hat{N}$ the particle number operator and $\tilde{\mu}$ the chemical potential. We write the change δE in the energy when the system is excited from its ground state as the sum of a quasiparticle (Landau) contribution, δE_L , and a multipair contribution, δE_M :

$$\delta E = \delta E_L + \delta E_M. \quad (2)$$

The quasiparticle contribution to the energy may be expressed as the sum of two terms, the intrinsic kinetic and mutual interaction energies of the quasiparticles, and the energy of interaction with the magnetic field:

$$\delta E_L = \delta E_L^{\text{int}} + \delta E_L^{\mathcal{H}}. \quad (3)$$

For simplicity, let us begin by considering a system of one species of fermions with spin 1/2. The change in the kinetic and interaction energies of the system when the quasiparticle distribution $n_{\mathbf{p}\sigma}$ changes by an amount $\delta n_{\mathbf{p}\sigma}$ is given to second order by the standard expression [26, 27], which amounts to the statement that the quasiparticle contribution to the energy is given by

$$\begin{aligned} \delta E_L^{\text{int}} = & \text{Tr}_\sigma \int \frac{d^3 p}{(2\pi\hbar)^3} (\epsilon_{\mathbf{p}}^0 - \tilde{\mu}) \delta n_{\mathbf{p}}(\sigma) \\ & + \frac{1}{2} \text{Tr}_\sigma \text{Tr}_{\sigma'} \int \frac{d^3 p}{(2\pi\hbar)^3} \int \frac{d^3 p'}{(2\pi\hbar)^3} f_{\mathbf{p}\mathbf{p}'\sigma\sigma'} \delta n_{\mathbf{p}}(\sigma) \delta n_{\mathbf{p}'}(\sigma'), \end{aligned} \quad (4)$$

where $\epsilon_{\mathbf{p}}$ is the quasiparticle energy. Note that $\delta n_{\mathbf{p}}(\sigma)$ is a matrix in spin space. The quantity $f_{\mathbf{p}\mathbf{p}'\sigma\sigma'}$ is the Landau quasiparticle interaction, which we write in the form

$$f_{\mathbf{p}\mathbf{p}'\sigma\sigma'} = f_{\mathbf{p}\mathbf{p}'} + g_{\mathbf{p}\mathbf{p}'} \boldsymbol{\sigma} \cdot \boldsymbol{\sigma}' + f_{\mathbf{p}\mathbf{p}'\sigma\sigma'}^{\text{T}}, \quad (5)$$

where $f_{\mathbf{p}\mathbf{p}'}$ is the spin-averaged quasiparticle interaction, $g_{\mathbf{p}\mathbf{p}'}$ is the spin-exchange contribution and $f_{\mathbf{p}\mathbf{p}'\sigma\sigma'}^{\text{T}}$ is the tensor or, more generally, the non-central contribution to the interaction. We shall discuss the form of the tensor interaction, $f_{\mathbf{p}\mathbf{p}'\sigma\sigma'}^{\text{T}}$, in Sec. IV in the light of the new results of Schwenk and Friman [7]. To begin, we derive an expression for the magnetic susceptibility, for a completely general interaction that conserves the total spin and the total angular momentum of the quasiparticle-quasihole pair, and subsequently we shall discuss its specific form in detail.

Non-central forces can alter the effective charges of the quasiparticles. When calculating the energy of interaction of the quasiparticles with an external field it is therefore important to allow for the fact that the effective couplings (charges) of quasiparticles may have components which are not scalars under rotations of the momentum of the quasiparticle. For definiteness, we consider the case of response to a magnetic field. The change in quasiparticle energy due to the application of a magnetic field, $\mathcal{H}$, is given by

$$\delta E_L^{\mathcal{H}} = -\text{Tr}_\sigma \sum_{ij} \mu_{ij} \sigma_j \mathcal{H}_i \delta n_{\mathbf{p}}(\sigma) \frac{d^3 p}{(2\pi\hbar)^3}, \quad (6)$$

where μ_{ij} , the magnetic moment matrix, is

$$\mu_{ij} = \mu \delta_{ij} + \frac{3}{2} \mu_{\text{T}} \left(\frac{p_i p_j}{p^2} - \frac{\delta_{ij}}{3} \right) \quad (7)$$

where μ and μ_{T} are parameters. Let us now consider the change in the energy when the Fermi surface is distorted. The distortion is specified by the function

$$\nu(\hat{\mathbf{p}})_{\alpha\beta} = \mathbf{u}(\hat{\mathbf{p}}) \cdot \boldsymbol{\sigma}_{\alpha\beta} \quad (8)$$

which corresponds to the change in the Fermi momentum as a function of direction. At zero temperature, the corresponding change in the distribution function $\delta n_{\mathbf{p}}$ may

be expanded in powers of ν and to second order one has

$$(\delta n_{\mathbf{p}})_{\alpha\beta} = \delta(p - p_F)\nu(\hat{\mathbf{p}})_{\alpha\beta} - \frac{1}{2}\delta'(p - p_F)\sum_{\gamma}\nu(\hat{\mathbf{p}})_{\alpha\gamma}\nu(\hat{\mathbf{p}})_{\gamma\beta}. \quad (9)$$

It is advantageous to use spherical tensors so, following Refs. [5, 23], we write

$$\nu(\hat{\mathbf{p}}) = \sum_{\nu=-1}^1 (-)^{\nu} u^{\nu}(\hat{\mathbf{p}}) \sigma^{-\nu} \quad (10)$$

and expand u in spherical harmonics

$$u^{\nu} = \sum_{l,m} u_{lm}^{\nu} Y_{lm}(\hat{\mathbf{p}}). \quad (11)$$

We shall work with eigenstates of the total angular momentum of the quasiparticle-quasihole pair. The corresponding amplitudes are constructed by the transformation

$$C_{lJ}^M = \sum_{m\mu} (-)^{\mu} (lm1 - \mu|JM) u_{lm}^{\mu}, \quad (12)$$

where $(lm1 - \mu|JM)$ are Clebsch-Gordan coefficients.

The part of the quasiparticle energy not involving the external field, Eq. (4), has been calculated in Refs. [5, 23] to be

$$\delta E_L^{\text{int}} = \frac{p_F^3}{(2\pi\hbar)^3 m^*} \sum_{l'l'JM} C_{lJ}^M C_{l'J}^{M*} \langle l'J|A|lJ \rangle, \quad (13)$$

where the matrix elements in the new basis are given by

$$\langle l'J'|A|lJ \rangle = \sum_{m,\mu,m',\mu'} (J'M'|l'm'1 - \mu') \langle l'm'\mu'|A|lm\mu \rangle (lm1 - \mu|JM), \quad (14)$$

$\langle l'J'|A|lJ \rangle$ are the normalized matrix elements, and

$$A = 1 + \mathcal{F}, \quad (15)$$

the unit coming from the first term in Eq. (3), and $\mathcal{F} = N(0)f_{\mathbf{p}\mathbf{p}'\sigma\sigma'}$ comes from the interaction term. For a quasiparticle interaction containing only exchange-tensor terms, the matrix elements $\langle lJ|A|l'J \rangle$ have been calculated for the conventional parametrization for the exchange tensor interaction in Ref. [5].

If we take the magnetic field to be in the z -direction, $\mathcal{H} = \mathcal{H}_z$, we find that the energy due to the application of this external field is

$$\delta E_L^{\mathcal{H}} = \frac{2p_F^2\sqrt{4\pi}}{(2\pi\hbar)^3} \mathcal{H}_z \times \left[\mu u_{00}^0 + \frac{\mu_T}{\sqrt{5}} u_{20}^0 - \frac{3}{2} \frac{\mu_T}{\sqrt{15}} (u_{21}^{-1} + u_{2-1}^1) \right]. \quad (16)$$

Rewriting this in terms of the coefficients C_{lJ}^M and adding the intrinsic contribution to the energy leads to the following expression for the quasiparticle energy:

$$\delta E_L = \frac{p_F^3}{(2\pi\hbar)^3 m^*} \left[\sum_{l'l'JM} C_{lJ}^M C_{l'J}^{M*} \langle l'J|A|lJ \rangle - \frac{2m^*}{p_F} \sqrt{4\pi} \mathcal{H}_z \left(\mu C_{01}^0 - \frac{\mu_T}{\sqrt{2}} C_{21}^0 \right) \right].$$

At first sight, one may be surprised that there are no terms with $J = 3$. However, this follows immediately from the observation that the magnetic moment operator $\mu_{ij}\sigma_j$ transforms as an axial vector under simultaneous rotations in momentum space and spin space, and consequently it contains only components corresponding to total angular momentum $J = 1$.

By minimizing the quasiparticle contribution to the energy with respect to the coefficients C_{lJ}^M we find that all C_{lJ}^M are zero except the following two,

$$C_{01}^0 = \sqrt{4\pi} \mathcal{H}_z \frac{m^*}{p_F} \left(\mu \langle 21|A|21 \rangle + \frac{\mu_T}{\sqrt{2}} \langle 01|A|21 \rangle \right) \times \frac{1}{\langle 21|A|21 \rangle \langle 01|A|01 \rangle - |\langle 01|A|21 \rangle|^2} \quad (17)$$

and

$$C_{21}^0 = \sqrt{4\pi} \mathcal{H}_z \frac{m^*}{p_F} \left(-\mu \langle 01|A|21 \rangle - \frac{\mu_T}{\sqrt{2}} \langle 01|A|01 \rangle \right) \times \frac{1}{\langle 21|A|21 \rangle \langle 01|A|01 \rangle - |\langle 01|A|21 \rangle|^2}. \quad (18)$$

III. STATIC RESPONSE FUNCTIONS

The magnetic susceptibility is defined as the derivative of the magnetization with respect to the magnetic field at zero field,

$$\chi = \left. \frac{\partial M_i}{\partial \mathcal{H}_i} \right|_{\mathcal{H}=0}. \quad (19)$$

We write the magnetization, M , as the sum of two terms, one contribution from single quasiparticle-quasihole pairs, M_L , and another contribution from the excitations of multipair states, which we denote by M_M :

$$M = M_L + M_M. \quad (20)$$

The first term, the Landau term, may be calculated by the same methods as we used in the previous section, and in terms of the C_{lJ}^M it is given by:

$$M_L = \int \sum_j \frac{d^3p}{(2\pi\hbar)^3} \mu_{ij} \sigma_j \delta n_{\mathbf{p}}(\sigma) = \frac{2p_F^2\sqrt{4\pi}}{(2\pi\hbar)^3} \left[\mu C_{01}^0 - \frac{\mu_T}{\sqrt{2}} C_{21}^0 \right]. \quad (21)$$

By using Eqs. (17) and (18) for the C_{ij}^M , and the relation between the susceptibility and the magnetization, we obtain the following expression for the susceptibility:

$$\begin{aligned} \chi = & N(0) \left[\mu^2 \frac{\langle 21|A|21 \rangle}{\langle 21|A|21 \rangle \langle 01|A|01 \rangle - \langle 01|A|21 \rangle^2} \right. \\ & + \mu \mu_T \sqrt{2} \frac{\langle 01|A|21 \rangle}{\langle 21|A|21 \rangle \langle 01|A|01 \rangle - \langle 01|A|21 \rangle^2} \\ & + \frac{\mu_T^2}{2} \frac{\langle 01|A|01 \rangle}{\langle 21|A|21 \rangle \langle 01|A|01 \rangle - \langle 01|A|21 \rangle^2} \left. \right] \\ & + \chi_M, \end{aligned} \quad (22)$$

where χ_M is the contribution from transitions to multipair states. Here we have implicitly made the usual choice of phase for the states, in which case the matrix elements of A are real. If we take only the first term and set $\mu = \mu_0$ in Eq. (22), this result reduces to that of Ref. [23]. Quantitative estimates of the different effects will be described in Sec. V after calculation of the relevant matrix elements, but first we shall discuss the form of the tensor interaction.

IV. THE TENSOR INTERACTION

The explicit form of the tensor interaction including all terms allowed by invariance under simultaneous rotations in coordinate space and spin space is [7]

$$f_{\mathbf{p}\mathbf{p}'}^T \sigma \sigma' = h_{\mathbf{p}\mathbf{p}'} S(\hat{\mathbf{q}}) + k_{\mathbf{p}\mathbf{p}'} S(\hat{\mathbf{P}}) + l_{\mathbf{p}\mathbf{p}'} A(\hat{\mathbf{q}}, \hat{\mathbf{P}}). \quad (23)$$

Here

$$S(\hat{\mathbf{q}}) = 3\sigma \cdot \hat{\mathbf{q}} \sigma' \cdot \hat{\mathbf{q}} - \sigma \cdot \sigma' \quad (24)$$

is referred to as the exchange-tensor operator, $\mathbf{q} = \mathbf{p} - \mathbf{p}'$ being the momentum transfer. The operator in the second term, which is referred to as the center of mass (cm) tensor, is given by

$$S(\hat{\mathbf{P}}) = 3\sigma \cdot \hat{\mathbf{P}} \sigma' \cdot \hat{\mathbf{P}} - \sigma \cdot \sigma'. \quad (25)$$

It has the same form as the exchange-tensor, but with $\mathbf{q}$ replaced by the total momentum $\mathbf{P} = \mathbf{p} + \mathbf{p}'$. The final term, referred to as the cross-vector term, is given by

$$\begin{aligned} A(\hat{\mathbf{q}}, \hat{\mathbf{P}}) &= (\sigma \times \sigma') \cdot (\hat{\mathbf{q}} \times \hat{\mathbf{P}}) \\ &= \sigma \cdot \hat{\mathbf{P}} \sigma' \cdot \hat{\mathbf{q}} - \sigma \cdot \hat{\mathbf{q}} \sigma' \cdot \hat{\mathbf{P}}. \end{aligned} \quad (26)$$

The functions $h_{\mathbf{p}\mathbf{p}'}$, $k_{\mathbf{p}\mathbf{p}'}$ and $l_{\mathbf{p}\mathbf{p}'}$ may be expanded in Legendre polynomials of $\cos \theta$, where θ is the angle between $\mathbf{p}$ and $\mathbf{p}'$, e.g. for $h_{\mathbf{p}\mathbf{p}'}$,

$$h_{\mathbf{p}\mathbf{p}'} = \sum_{l=0}^{\infty} h_l P_l(\cos \theta). \quad (27)$$

Note that the choice of parametrization of the exchange-tensor term in Eq. (23) is different than the one traditionally used. The conventional way to parametrize this term is to write [2]

$$h_{\mathbf{p}\mathbf{p}'} S(\hat{\mathbf{q}}) = \frac{q^2}{p_F^2} \tilde{h}_{\mathbf{p}\mathbf{p}'} S(\hat{\mathbf{q}}). \quad (28)$$

The functions h and $\tilde{h}$ contain the same physical information, but have different properties as far as their expansions in terms of Legendre polynomials are concerned. We will now compare some aspects of the different choices of parametrization.

With the conventional parametrization, $\tilde{h}_{\mathbf{p}\mathbf{p}'}$, the explicit q^2 factor ensures that the tensor interaction vanishes identically for $\mathbf{q} \rightarrow 0$, provided that $\tilde{h}_{\mathbf{p}\mathbf{p}'}$ is finite for $\mathbf{p} \rightarrow \mathbf{p}'$. This is in agreement with the form of the one-pion-exchange contribution which is given by

$$f_{\mathbf{p}\mathbf{p}'\sigma\sigma'}^T = \frac{f^2}{3m_\pi^2} \frac{q^2}{q^2 + m_\pi^2} S(\hat{\mathbf{q}}), \quad (29)$$

where f is the pion-nucleon coupling constant, and $f^2/(4\pi\hbar c) \approx 0.08$. More generally, if the tensor interaction is analytic for small $\mathbf{q}$, one would expect $h_{\mathbf{p}\mathbf{p}'}$ to tend to zero for $\mathbf{p} \rightarrow \mathbf{p}'$, since otherwise the tensor interaction would depend on the way in which $\mathbf{p}$ approaches $\mathbf{p}'$. Such nonanalytic contributions to the quasiparticle interaction do occur for systems with central interactions [28, 29], and there they are due to exchange of quasiparticle-quasihole pairs and collective modes with long wavelengths. The work of Schwenk and Friman [7] indicates that such contributions do not occur in $h_{\mathbf{p}\mathbf{p}'}$, although they do in $k_{\mathbf{p}\mathbf{p}'}$ (see Sec. VB). To ensure that $h_{\mathbf{p}\mathbf{p}'} \rightarrow 0$ for $\mathbf{p} \rightarrow \mathbf{p}'$, the h_l must satisfy the condition

$$\sum_{l=0}^{\infty} h_l = 0. \quad (30)$$

If the series is truncated after a finite number of terms, this will inevitably lead to the sum rule being violated. However, this should not be very important as long as long-wavelength processes do not play a dominant role.

The conventional parametrization suffers from the disadvantage that, when expanding $\tilde{h}$ in terms of Legendre polynomials (27), it is generally necessary to include many terms in the sum in order to obtain a good representation of the interaction. Consider, for example, the one pion exchange contribution. Using the fact that $\cos \theta = 1 - q^2/(2p_F^2)$ one finds

$$\begin{aligned} \tilde{h}_l^{\text{OPE}} &= \\ &= (2l+1) \frac{f^2}{(3m_\pi^2)} \int_0^{2p_F} \frac{dq^2}{p_F^2} \frac{1}{q^2 + m_\pi^2} P_l[1 - q^2/(2p_F^2)] \\ &= (2l+1) \frac{f^2}{(3m_\pi^2)} Q_l[1 + \frac{m_\pi^2}{2p_F^2}], \end{aligned} \quad (31)$$

where Q_l is a Legendre function of the second kind. In

TABLE I: Tensor parameters for neutron matter with Fermi wave vector $k_F = 1.7 \text{ F}^{-1}$. Values for the tensor parameters by Schwenk [30], obtained taking into account the tensor interaction to first order by the renormalization group method described in Refs. [6, 31], are presented in the first and third columns, for the two choices of parametrization. For comparison we present the result for the one-pion exchange potential (Eq. (32)) in column two and four. All tensor parameters are calculated for $m^*/m = 0.8345$.

l	H_l	H_l^{OPE}	$\tilde{H}_l$	$\tilde{H}_l^{\text{OPE}}$
0	0.529	0.403	0.665	0.751
1	0.150	-0.088	1.200	1.037
2	-0.0959	-0.079	1.150	0.935
3	-0.141	-0.064	0.933	0.753
4	-0.124	-0.049	0.703	0.575
5	-0.0944	-0.036	0.509	0.427
6	-0.0697	-0.026	0.359	0.310
7	-0.0494	-0.019	0.249	0.223
8	-0.0335	-0.013	0.171	0.158
9	-0.0225	-0.009	0.116	0.111
10	-0.0151	-0.007	0.0774	0.078

Table 1 we show results of calculations of the exchange tensor interaction, $\tilde{H}_l = N(0)\tilde{h}_l$, by Schwenk [30], for $k_F = 1.7 \text{ F}^{-1}$. Note that these were calculated taking the tensor interaction into account only in first order. One sees that as l increases from zero, $\tilde{H}_l$ first increases and then decreases. It is therefore a poor approximation to truncate the expansion after the first few terms. The reason for this is that typical Fermi wave numbers at nuclear densities are of order 1.5 F^{-1} , which is large compared with the pion mass, $m_\pi/\hbar c \sim 0.7 \text{ F}^{-1}$. In the limit of the zero pion mass, all the $\tilde{h}_l$ diverge, the leading contribution being $(2l+1)f^2/(3m_\pi^2)\ln(2p_F/m_\pi)$. This is due to the fact that $\tilde{h} \sim q^{-2}$, which leads to a logarithmic integral for $\tilde{h}_l$.

The situation is quite different for the parametrization (27). For the one pion exchange potential (29) one finds

$$h_l = \frac{f^2}{(3m_\pi^2)} \left(\delta_{l,0} - \frac{m_\pi^2}{2p_F^2} (2l+1)Q_l[1 + m_\pi^2/2p_F^2] \right). \quad (32)$$

This shows that for small m_π/p_F , the leading term is

$$h_l \sim \frac{f^2}{3m_\pi^2} \frac{2l+1}{2} \left(\frac{m_\pi}{p_F} \right)^2 \ln(2p_F/m_\pi). \quad (33)$$

In the limit of zero pion mass, $h_{\mathbf{pp}'}$ becomes a constant, and therefore the only nonvanishing coefficient in the expansion in Legendre polynomial is h_0 . Note that for both $\tilde{h}_l$ and h_l , the limits $m_\pi/p_F \rightarrow 0$ and $l \rightarrow \infty$ do not commute.

Let us now compare the expansions of h_l and $\tilde{h}_l$. Using the relation between $h_{\mathbf{pp}'}$ and $\tilde{h}_{\mathbf{pp}'}$, Eq. (28) and that $1 - \cos\theta = q^2/(2p_F^2)$, we find the following expression for

h_l :

$$h_l = 2(2l+1) \int_{-1}^{+1} \frac{d(\cos\theta)}{2} (1 - \cos\theta) \tilde{h}(\cos\theta) P_l(\cos\theta) \quad (34)$$

or

$$h_l = 2 \left(\tilde{h}_l - \frac{l+1}{2l+3} \tilde{h}_{l+1} - \frac{l}{2l-1} \tilde{h}_{l-1} \right). \quad (35)$$

Thus, given the $\tilde{h}_l$'s, we can obtain the h_l 's. Next we show how to write the $\tilde{h}_l$'s, given by

$$\tilde{h}_l = \frac{(2l+1)}{2} \int_{-1}^1 \frac{d(\cos\theta)}{2} \frac{h(\cos\theta)}{1 - \cos\theta} P_l(\cos\theta) \quad (36)$$

in terms of the h_l 's. Expanding the function $h(\cos\theta)$ in Legendre polynomials, we get

$$\tilde{h}_l = \frac{(2l+1)}{2} \int_{-1}^1 \frac{d(\cos\theta)}{2} \sum_{l'} h_{l'} P_{l'}(\cos\theta) \frac{P_l(\cos\theta)}{1 - \cos\theta}. \quad (37)$$

Since the series on the right hand side is not uniformly convergent for $\cos\theta$ in the closed interval $[-1, 1]$, we cannot invert the order of the sum and the integral. However, because of the sum rule (30), $h(\theta=0)$ vanishes, there is no divergence, and we can rewrite Eq. (36) as

$$\tilde{h}_l = \frac{(2l+1)}{2} \int_{-1}^1 \frac{d(\cos\theta)}{2} \frac{(h(\cos\theta) - h(\theta=0))}{1 - \cos\theta} P_l(\cos\theta). \quad (38)$$

Making use of the sum rule, we find that

$$\tilde{h}_l = \frac{(2l+1)}{2} \int_{-1}^1 \frac{d(\cos\theta)}{2} \sum_{l'} h_{l'} [P_{l'}(\cos\theta) - 1] \frac{P_l(\cos\theta)}{1 - \cos\theta}. \quad (39)$$

This expression may be evaluated using a number of standard results. First

$$\int_{-1}^1 \frac{d(\cos\theta)}{2} \frac{[P_{l'}(\cos\theta) - P_0(z)]}{z - \cos\theta} P_l(\cos\theta) = P_{l'}(z)Q_l(z) - P_0(z)Q_l(z), \quad l' \leq l. \quad (40)$$

The $Q_l(z)$ may be expressed as

$$Q_l(z) = \frac{1}{2} P_l(z) \ln \frac{1+z}{1-z} - W_{l-1}, \quad (41)$$

where

$$W_{l-1} = \sum_{k=1}^l \frac{1}{k} P_{k-1}(z) P_{l-k}(z) \quad \text{and} \quad (42)$$

$$W_{-1} = 0. \quad (43)$$

The Legendre polynomial $P_l(z)$ may be written in the following form:

$$P_l(z) = 2^{-l} \sum_{k=1}^{[l/2]} (-1)^l \binom{l}{k} \binom{2l-2k}{n} z^{l-2k}. \quad (44)$$

Letting $z \rightarrow 1$ in the integral, we find that the logarithmic divergence vanishes and the final result is

$$\tilde{h}_l = -\frac{(2l+1)}{2} \sum_{l'=l+1}^{\infty} h_{l'} \sum_{k=l+1}^{l'} \frac{1}{k}. \quad (45)$$

V. EVALUATION OF THE MATRIX ELEMENTS

We return to the calculation of the matrix elements of $A = 1 + \mathcal{F}$, Eq. (15), which we write as

$$\langle lJ|A|l'J \rangle = \delta_{ll'} + \langle lJ|\mathcal{F}|l'J \rangle. \quad (46)$$

We further write $\langle lJ|\mathcal{F}|l'J \rangle$ as the sum of a number of contributions:

$$\begin{aligned} \langle lJ|\mathcal{F}|l'J \rangle &= \langle lJ|\mathcal{F}^c|l'J \rangle + \langle lJ|\mathcal{F}^{\text{ex}}|l'J \rangle \\ &+ \langle lJ|\mathcal{F}^{\text{cm}}|l'J \rangle + \langle lJ|\mathcal{F}^{\text{cv}}|l'J \rangle. \end{aligned} \quad (47)$$

Here $\mathcal{F}^c$ is the central part, $\mathcal{F}^{\text{ex}}$ is the exchange tensor part, $\mathcal{F}^{\text{cm}}$ is the center of mass tensor part and $\mathcal{F}^{\text{cv}}$ is the cross vector part.

A. Central and exchange tensor contributions

In Ref. [5] the central and exchange tensor part in the conventional parametrization were calculated to be

$$\langle lJ|\mathcal{F}^c|l'J \rangle = \delta_{ll'} \frac{G_l}{2l+1}, \quad (48)$$

and

$$\begin{aligned} \langle lJ|\mathcal{F}^{\text{ex}}|l'J \rangle &= \delta_{ll'} \left(\tilde{H}_{l-1} \frac{2l}{(2l-1)(2l+1)} + \tilde{H}_{l+1} \frac{2(l+1)}{(2l+3)(2l+1)} \right) \\ &+ (-1)^J 15 \begin{pmatrix} 1 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l & l' & 2 \\ 0 & 0 & 0 \end{pmatrix} \left\{ \begin{matrix} 1 & l' & J \\ l & 1 & 2 \end{matrix} \right\} \left[\tilde{H}_l \left(\frac{2l+1}{2l'+1} \right)^{1/2} + \tilde{H}_{l'} \left(\frac{2l'+1}{2l+1} \right)^{1/2} \right] \\ &- \frac{3[(2l+1)(2l'+1)]^{1/2}}{2J+1} \begin{pmatrix} l & 1 & J \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l' & 1 & J \\ 0 & 0 & 0 \end{pmatrix} \tilde{H}_J - 3[(2l+1)(2l'+1)]^{1/2} \sum_{l''} \begin{pmatrix} l & 1 & l'' \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l' & 1 & l'' \\ 0 & 0 & 0 \end{pmatrix} \left\{ \begin{matrix} 1 & l' & J \\ 1 & l & l'' \end{matrix} \right\} \tilde{H}_{l''}. \end{aligned} \quad (49)$$

Here $\begin{pmatrix} l_1 & l_2 & l \\ m_1 & m_2 & m \end{pmatrix}$ are Wigner $3j$ -symbols and

$\left\{ \begin{matrix} l_1 & l_2 & l_3 \\ l'_1 & l'_2 & l'_3 \end{matrix} \right\}$ are Wigner $6j$ -symbols.

We can transform the matrix elements above to the new parametrization. In the expression (22) we can see that the only matrix elements that are necessary for the calculation of the susceptibility are $\langle 01|A|01 \rangle$, $\langle 01|A|21 \rangle$ and $\langle 21|A|21 \rangle$. The contribution to these matrix elements from non-interacting quasiparticles and from the central part of the quasiparticle interaction is given by

$$\langle l'J|1 + \mathcal{F}^c|lJ \rangle = \delta_{ll'} \left(1 + \frac{G_l}{2l+1} \right) \quad (50)$$

and the exchange tensor contributions by [5]

$$\begin{aligned} \langle 01|\mathcal{F}^{\text{ex}}|01 \rangle &= 0, \\ \langle 01|\mathcal{F}^{\text{ex}}|21 \rangle &= -\sqrt{2} \left(\tilde{H}_0 - \frac{2}{3} \tilde{H}_1 + \frac{1}{5} \tilde{H}_2 \right) \text{ and} \end{aligned}$$

$$\langle 21|\mathcal{F}^{\text{ex}}|21 \rangle = -\frac{7}{15} \tilde{H}_1 + \frac{2}{5} \tilde{H}_2 - \frac{3}{35} \tilde{H}_3. \quad (51)$$

Let us now express this result in terms of H_l . As an example we consider the combination $\tilde{H}_0 - 2\tilde{H}_1/3 + \tilde{H}_2/5$. This may be written in the form

$$\begin{aligned} &\tilde{H}_0 - \frac{2}{3} \tilde{H}_1 + \frac{1}{5} \tilde{H}_2 \\ &= \int_{-1}^{+1} \frac{d(\cos \theta)}{2} \tilde{H}(\cos \theta) [P_0(\cos \theta) - 2P_1(\cos \theta) + P_2(\cos \theta)] \\ &= \int_{-1}^{+1} \frac{d(\cos \theta)}{2} H(\cos \theta) \frac{[P_0(\cos \theta) - 2P_1(\cos \theta) + P_2(\cos \theta)]}{2(1 - \cos \theta)}. \end{aligned} \quad (52)$$

An interesting property of the fraction in Eq. (52) is that it is finite for $\theta \rightarrow 0$, and therefore the integral converges if $H(\cos \theta)$ is finite in this limit, even though individual terms diverge. One finds similar results for the other combinations of the $\tilde{H}_l$ that occur in Eqs. (51). The

corresponding result written in terms of the H_l are

$$\begin{aligned}\langle 01|\mathcal{F}^{\text{ex}}|01\rangle &= 0, \\ \langle 01|\mathcal{F}^{\text{ex}}|21\rangle &= -\frac{1}{2\sqrt{2}}(H_0 - H_1) \quad \text{and} \\ \langle 21|\mathcal{F}^{\text{ex}}|21\rangle &= \frac{H_2}{10} - \frac{1}{4}(H_0 + H_1).\end{aligned}\quad (53)$$

It remains to calculate the center of mass tensor and the cross vector contributions.

B. Center of mass tensor

The calculation of the center of mass tensor terms follows closely that of the exchange tensor. Let us study the operator $\mathcal{F}^{\text{cm}}$ given by

$$\mathcal{F}_{\mathbf{p}\mathbf{p}'}^{\text{cm}} = N(0)k_{\mathbf{p}\mathbf{p}'}S(P) = \sum_l K_l P_l(\cos\theta)S(P). \quad (54)$$

If we reverse one of the momenta, we find that

$$\begin{aligned}\mathcal{F}_{\mathbf{p}-\mathbf{p}'}^{\text{cm}} &= N(0)k_{\mathbf{p}-\mathbf{p}'}S(q) \\ &= \sum_l (-1)^l K_l P_l(\cos\theta)S(q).\end{aligned}\quad (55)$$

Thus we have a similar expression to the exchange tensor operator, with a difference of a factor of $(-1)^l$. For the state vectors $|l'J'\rangle$, reversing the momenta, gives

$$\langle -\hat{\mathbf{p}}'\sigma'|l'J'\rangle = (-1)^{l'}\langle \hat{\mathbf{p}}'\sigma'|l'J'\rangle. \quad (56)$$

The matrix elements of $\mathcal{F}^{\text{cm}}$ is thus the same as those for $\mathcal{F}^{\text{ex}}$, apart from an overall factor of $(-1)^{l'}$ and that H_l should be replaced by $(-1)^l K_l$.

However, when calculating the matrix elements of the center of mass tensor, it is necessary to be careful for $\theta \rightarrow \pi$, since the calculation of $k_{\mathbf{p}\mathbf{p}'}$ by Schwenk and Friman [7] showed that $k_{\mathbf{p}\mathbf{p}'}$ does not vanish in this limit, but rather tends to a constant when second order tensor contributions are included. However, we shall show, that for the matrix elements we consider, this will not pose a problem.

By writing $k_{\mathbf{p}\mathbf{p}'}$ as $(P^2/p_F^2)\tilde{k}_{\mathbf{p}\mathbf{p}'}$ we can use the calculation for the exchange tensor given in Ref. [5], except that for the odd l' states we have to include a factor $(-1)^{l'}$, and $\tilde{H}_l$ should be replaced by $(-1)^l \tilde{K}_l$, with $\tilde{K}_l = N(0)\tilde{k}_l$. For the particular matrix elements that we are interested in we find that

$$\begin{aligned}\langle 01|\mathcal{F}^{\text{cm}}|21\rangle &= -\sqrt{2}\left(\tilde{K}_0 + \frac{2}{3}\tilde{K}_1 + \frac{1}{5}\tilde{K}_2\right) \quad \text{and} \\ \langle 21|\mathcal{F}^{\text{cm}}|21\rangle &= \frac{7}{15}\tilde{K}_1 + \frac{2}{5}\tilde{K}_2 + \frac{3}{35}\tilde{K}_3.\end{aligned}\quad (57)$$

Just as in the example (52), we find that the matrix elements above give rise to combinations of Legendre polynomials which contain a factor $1 + \cos\theta$, so the integral

will be finite for $\theta \rightarrow \pi$, provided that $\tilde{k}_{\mathbf{p}\mathbf{p}'}$ is finite in this limit. Using the result of Schwenk and Friman [7] that $k_{\mathbf{p}\mathbf{p}'}$ goes to a constant when $\theta \rightarrow \pi$, we can see that $\tilde{k}_l$, which is given by

$$\tilde{k}_l = \frac{2l+1}{2p_F^2} \int_{-1}^1 \frac{k_{\mathbf{p}\mathbf{p}'}}{1+\cos\theta} P_l(\cos\theta) d(\cos\theta), \quad (58)$$

will, in fact, have a logarithmic divergence. To deal with this divergence, we replace the limit -1 by Λ , and write $\tilde{k}_l$ as the sum of a regular part and a part that contains the logarithmic divergence,

$$\tilde{k}_l(\Lambda) = \tilde{k}_l^{\text{reg}} + (-1)^l(2l+1)\ln(1+\Lambda)C(\Lambda) \quad (59)$$

where $C(\Lambda) = k_{\mathbf{p}\mathbf{p}'}(\Lambda)/(2p_F^2)$ tends to a constant when $\Lambda \rightarrow -1$ and the factor $(-1)^l$ comes from $P_l(-1) = (-1)^l$. By inserting the expression above in the matrix elements of $\mathcal{F}^{\text{cm}}$, Eq. (57), we obtain

$$\begin{aligned}\langle 01|\mathcal{F}^{\text{cm}}|21\rangle &= -\sqrt{2}\left(\tilde{K}_0^{\text{reg}} + C(\Lambda)\ln(1+\Lambda)\right. \\ &\quad + \frac{2}{3}\tilde{K}_1^{\text{reg}} - 2C(\Lambda)\ln(1+\Lambda) \\ &\quad + \left.\frac{1}{5}\tilde{K}_2^{\text{reg}} + C(\Lambda)\ln(1+\Lambda)\right) \\ &= -\sqrt{2}\left(\tilde{K}_0^{\text{reg}} + \frac{2}{3}\tilde{K}_1^{\text{reg}} + \frac{1}{5}\tilde{K}_2^{\text{reg}}\right) \quad (60) \\ \text{and} \\ \langle 21|\mathcal{F}^{\text{cm}}|21\rangle &= \frac{7}{15}[\tilde{K}_1^{\text{reg}} - 3C(\Lambda)\ln(1+\Lambda)] \\ &\quad + \frac{2}{5}\tilde{K}_2^{\text{reg}} + 2C(\Lambda)\ln(1+\Lambda) \\ &\quad + \frac{3}{35}[\tilde{K}_3^{\text{reg}} - 7C(\Lambda)\ln(1+\Lambda)] \\ &= \frac{7}{15}\tilde{K}_1^{\text{reg}} + \frac{2}{5}\tilde{K}_2^{\text{reg}} + \frac{3}{35}\tilde{K}_3^{\text{reg}}.\end{aligned}\quad (61)$$

This shows that in the combinations of parameters we need, the divergent contribution vanishes and therefore allows us to simply replace H_l by $(-1)^l K_l$ in the expression (53) for the $\langle lJ|\mathcal{F}^{\text{ex}}|l'J\rangle$ to obtain $\langle lJ|\mathcal{F}^{\text{cm}}|l'J\rangle$ in terms of the K_l 's:

$$\begin{aligned}\langle 01|\mathcal{F}^{\text{cm}}|01\rangle &= 0, \\ \langle 01|\mathcal{F}^{\text{cm}}|21\rangle &= -\frac{1}{2\sqrt{2}}(K_0 + K_1) \quad \text{and} \\ \langle 21|\mathcal{F}^{\text{cm}}|21\rangle &= \frac{K_2}{10} - \frac{1}{4}(K_0 - K_1).\end{aligned}\quad (62)$$

C. The cross vector term

Finally, we turn to the cross vector term, which is proportional to

$$A(\hat{\mathbf{q}}, \hat{\mathbf{P}}) = (\boldsymbol{\sigma} \times \boldsymbol{\sigma}') \cdot (\hat{\mathbf{q}} \times \hat{\mathbf{P}})$$

$$\begin{aligned}
&= (\boldsymbol{\sigma} \times \boldsymbol{\sigma}') \cdot \left(\frac{\hat{\mathbf{p}} \times \hat{\mathbf{p}}'}{|\hat{\mathbf{p}} \times \hat{\mathbf{p}}'|} \right) \\
&= \frac{\boldsymbol{\sigma} \cdot \hat{\mathbf{p}} \boldsymbol{\sigma}' \cdot \hat{\mathbf{p}}' - \boldsymbol{\sigma} \cdot \hat{\mathbf{p}}' \boldsymbol{\sigma}' \cdot \hat{\mathbf{p}}}{|\hat{\mathbf{p}} \times \hat{\mathbf{p}}'|}. \quad (63)
\end{aligned}$$

The calculations of Ref. [7] indicate that this term will be well behaved in the limits where $\sin \theta \rightarrow 0$, since $L_{\mathbf{pp}'} \rightarrow 0$ for both $\theta = 0$ and $\theta = \pi$. This means that there will be sum rules like the one (30) for both $\theta = 0$ and $\theta = \pi$:

$$\begin{aligned}
\sum_l L_l P_l(\cos \theta = 1) &= 0, \text{ and} \\
\sum_l L_l P_l(\cos \theta = -1) &= \sum_l L_l (-1)^l P_l(\cos \theta = 1) = 0
\end{aligned}$$

which gives two sum rules, one for the odd coefficients and one for even ones:

$$\sum_{l=0}^{\infty} L_{2l} = 0, \text{ and}$$

$$\sum_{l=0}^{\infty} L_{2l+1} = 0. \quad (64)$$

To simplify the calculation of the matrix elements, we define $\tilde{L}_{\mathbf{pp}'} = L_{\mathbf{pp}'} / \sin \theta$, and calculate the matrix element $\langle lJ | \mathcal{F}^{\text{cv}} | l'J \rangle$, which later will be transformed back to the original notation. The cross vector contribution to the quasiparticle interaction is

$$\Delta \mathcal{F}^{\text{cv}} = N(0) \frac{1}{4\pi} \frac{1}{4} \text{Tr} \boldsymbol{\sigma} \boldsymbol{\sigma}' \iint d\Omega d\Omega' \mathbf{u} \cdot \boldsymbol{\sigma} \mathcal{F}^{\text{cv}} \mathbf{u}' \cdot \boldsymbol{\sigma}'. \quad (65)$$

We use the notation and method described in Sec. II, which gives for the matrix elements in the $|lm1-\mu\rangle$ basis

$$\langle lm1-\mu | \mathcal{F}^{\text{cv}} | l'm'1-\mu' \rangle = \sum_{l'',m''} (-)^{m''} \tilde{L}_{l''} \times \quad (66)$$

$$\left[\begin{pmatrix} l & 1 & l'' \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l' & 1 & l'' \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l & 1 & l'' \\ m & -\mu & m'' \end{pmatrix} \begin{pmatrix} l' & 1 & l'' \\ m' & -\mu' & m'' \end{pmatrix} - \begin{pmatrix} l & 1 & l'' \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l' & 1 & l'' \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l & 1 & l'' \\ m & -\mu' & m'' \end{pmatrix} \begin{pmatrix} l' & 1 & l'' \\ m' & -\mu & m'' \end{pmatrix} \right].$$

Changing to the $|lJ\rangle$ basis, we find the matrix element to be

$$\begin{aligned}
&\langle lJ | \mathcal{F}^{\text{cv}} | l'J \rangle = \\
&+ \frac{[(2l+1)(2l'+1)]^{1/2}}{2J+1} \begin{pmatrix} l & 1 & J \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l' & 1 & J \\ 0 & 0 & 0 \end{pmatrix} \tilde{L}_J \\
&- [(2l+1)(2l'+1)]^{1/2} \\
&\sum_{l''} \begin{pmatrix} l & 1 & l'' \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l' & 1 & l'' \\ 0 & 0 & 0 \end{pmatrix} \begin{Bmatrix} 1 & l' & J \\ 1 & l & l'' \end{Bmatrix} \tilde{L}_{l''}. \quad (67)
\end{aligned}$$

For $l' = l + 2$, the triangle conditions imply that the 3-j symbols are non-zero only if $J = l + 1$ and $l'' = l + 1$. When this happens the two terms are equal in magnitude but of opposite sign. Therefore all $\langle lJ | \mathcal{F}^{\text{cv}} | (l+2)J \rangle$ are zero. The only non-zero matrix elements are when $l = l'$. Then J can be either $l + 1$ or $l - 1$:

$$\begin{aligned}
&\langle lJ | \mathcal{F}^{\text{cv}} | lJ \rangle = \\
&+ \frac{2(l+1)^2}{(2l+3)^2(2l+2)} \left[\frac{1}{(l+1)(2l+1)-1} \right] \tilde{L}_{l+1} \\
&- \frac{l}{(2l+1)(2l-1)} \tilde{L}_{l-1}, \text{ for } J = l + 1, \text{ and} \quad (68) \\
&\langle lJ | \mathcal{F}^{\text{cv}} | lJ \rangle =
\end{aligned}$$

$$\begin{aligned}
&+ \frac{l}{(2l-1)^2} \left[\frac{1}{l(2l+1)-1} \right] \tilde{L}_{l-1} \\
&- \frac{2(l+1)^2}{(2l+1)(2l+2)(2l+3)} \tilde{L}_{l+1}, \text{ for } J = l - 1 \quad (69)
\end{aligned}$$

The matrix elements we need thus have the following contributions from the cross vector:

$$\begin{aligned}
\langle 01 | \mathcal{F}^{\text{cv}} | 01 \rangle &= 0, \\
\langle 21 | \mathcal{F}^{\text{cv}} | 01 \rangle &= 0 \quad \text{and} \\
\langle 21 | \mathcal{F}^{\text{cv}} | 21 \rangle &= \frac{2}{10} \tilde{L}_1 - \frac{3}{35} \tilde{L}_3. \quad (70)
\end{aligned}$$

The only matrix element we need for the cross vector term is thus when $l = l' = 2$ and $J = 1$, and we transform this to our original parametrization by writing:

$$\begin{aligned}
\langle 21 | \mathcal{F}^{\text{cv}} | 21 \rangle &= \quad (71) \\
&\int_{-1}^1 d(\cos \theta) \frac{3}{5} \sum_k \frac{L_k P_k(\cos \theta)}{\sin \theta} [P_1(\cos \theta) - P_3(\cos \theta)]
\end{aligned}$$

which is the integral

$$-\frac{3}{2} \int_{-1}^1 dx \sum_n L_n P_n(x) \sqrt{1-x^2} x = -\frac{3}{2} \sum_n L_n I_n. \quad (72)$$

TABLE II: Landau parameters for neutron matter calculated by Schwenk [7]. The Landau parameters are calculated for Fermi wave vector $k_F = 1.7 \text{ F}^{-1}$ and the effective mass is $m^*/m = 0.8345$.

l	G_l	H_l	K_l	L_l
0	0.842	0.070	-0.258	-0.060
1	0.412	0.163	0.146	-0.089
2	0.219	-0.301	0.124	-0.034
3	0.109	-0.002	0.048	0.025
4	0.051	-0.150	0.015	0.043

We calculate the integral I_n to be

$$I_n = \begin{cases} \sum_{k=0}^{[n/2]} \frac{\pi(-1)^k (2n-2k)!(n-2k+1)}{k!(n-k)!(\frac{n-2k+1}{2})!(\frac{n-2k+3}{2})!}, & n = 2l + 1 \\ 0 & n = 2l \end{cases} \quad (73)$$

D. Matrix elements and the susceptibility

Combining Eqs. (50),(53),(62),(72) and (72) above, we find

$$\begin{aligned} \langle 01|A|01 \rangle &= 1 + G_0, \\ \langle 01|A|21 \rangle &= -\frac{1}{2\sqrt{2}} [H_0 - H_1 + K_0 + K_1] \quad \text{and} \\ \langle 21|A|21 \rangle &= 1 + \frac{1}{5} \left(G_2 + \frac{H_2 + K_2}{2} \right) \\ &\quad - \frac{1}{4} (H_0 + K_0 + H_1 - K_1) - \frac{3}{2} \sum_n L_n I_n, \end{aligned} \quad (74)$$

with I_n given by Eq. (73).

From these results one may calculate the magnetic susceptibility from Eq. (22). The effects of the tensor interaction are generally small, so to estimate the various contributions to the susceptibility we expand the susceptibility to second order in the tensor interaction and μ_T and find

$$\begin{aligned} \chi &= N(0) \left[\mu^2 \frac{1}{1+G_0} + \mu^2 \frac{1}{(1+G_0)^2} \frac{\langle 01|A|21 \rangle^2}{1+G_2/5} \right. \\ &\quad + \mu \mu_T \sqrt{2} \frac{\langle 01|A|21 \rangle}{(1+G_0)(1+G_2/5)} \\ &\quad \left. + \frac{\mu_T^2}{2} \frac{1}{1+G_2/5} \right] + \chi_M. \end{aligned} \quad (75)$$

It is not easy to obtain a reliable estimate of the various terms, since different calculations of the tensor Landau parameters give different result. To exemplify this for

pure neutrons, we first use the Landau parameters obtained by Schwenk [30] listed Table I. These take into account the tensor interaction only to first order, and consequently the only tensor term is the exchange tensor. We find

$$\begin{aligned} \langle 01|A|21 \rangle &\approx -0.13 \quad \text{and} \\ \langle 21|A|21 \rangle &\approx 0.86. \end{aligned} \quad (76)$$

We can now compare this with what we get if we use the Landau parameters of Ref. [7], listed in Table II. These were calculated including the tensor interaction to second order. We find that the exchange tensor, center of mass tensor and the cross vector terms are all quite small, so the matrix elements are only slightly modified from the value one obtains using only the central part of the interaction. In addition to being small, the tensor contributions have a different sign from the first order result:

$$\begin{aligned} \langle 01|A|21 \rangle &\approx 0.07 \quad \text{and} \\ \langle 21|A|21 \rangle &\approx 1.06 \quad \text{with} \\ \langle 21|1 + \mathcal{F}^c|21 \rangle &\approx 1.04. \end{aligned} \quad (77)$$

Since the calculation of tensor parameters in Ref. [7] is only to second order, and the second order effects almost completely cancel the first order ones, higher order effects could be significant and should be calculated.

From the two sets of results above we can draw some conclusions regarding the effect of the tensor interaction on the susceptibility. The second term in Eq. (75) is clearly small compared to the first term, of the order of 1%, and may be neglected. The third and fourth terms, could be important, depending on the value of the tensor magnetic moment. The corrections to the bare nucleon moment arise from two sources. The first is configuration mixing, which is due to the fact that a quasiparticle in the medium consists of a bare nucleon and a superposition of more complicated states involving a nucleon plus a number of particle-hole pairs. The second source is exchange currents, which are due to the fact that the external field can interact with other degrees of freedom than nucleons, for example, intermediate mesons in a nucleon-nucleon interaction. For definiteness, we shall use the results given by Arima *et al.* in Ref. [21] for the Fermi gas model. For neutrons in symmetric nuclear matter, they find $\mu \approx -1.62 \mu_N$ and $\mu_T \approx 0.08 \mu_N$, where $\mu_N = e\hbar/2M$ is the nuclear magneton. Calculations of the configuration mixing contribution to weak interaction matrix elements made by Cowell and Pandharipande [32] for asymmetric nuclear matter indicate that the renormalization of the magnetic moment does not depend strongly on proton fraction, and amounts to a suppression by about 10%. We would thus suspect that renormalization of the magnetic moment would be relatively insensitive to the proton fraction, and would be of a similar order of magnitude. However, we stress that it is important to make more detailed estimates of magnetic moments in the nuclear medium.

The small magnitude of the tensor magnetic moment means that we may neglect also the third and fourth term in Eq. (75), both being less than one per cent of the first term. Therefore, we conclude that, the most important effects of the tensor force on the susceptibility is most likely the renormalization of the isotropic part of the magnetic moment and to the presence of transitions to multipair states.

No reliable calculation of χ_M exists, but by using the sum rule argument presented in Ref. [19] we can put a lower bound on the multipair contribution to the susceptibility:

$$\chi_M(q, 0) \geq \frac{2nS(q)}{\bar{\omega}}, \quad (q \rightarrow 0). \quad (78)$$

Here $S(q)$ is the static structure function, n is the density and $\bar{\omega}$ is the mean excitation energy. In Ref. [19] we took the values of $S(q) = 0.19$ and $\bar{\omega} = 63$ MeV from the calculations of Ref. [33] and estimated the multipair contribution to the susceptibility to be more than 60% of the susceptibility calculated by Fantoni et al [18] ($\chi = 0.38\chi_F$). Here χ_F is the susceptibility of a free Fermi gas of neutrons, $\chi_F = 3n/2\epsilon_F$. New calculations of the static structure function and the mean excitation energy by Cowell and Pandharipande [34] give a similar result, $S(q = 0) = 0.187$ and $\bar{\omega} = 64$ MeV. In order to compare with symmetric nuclear matter in next section we calculate χ_M/χ_F , and we find it to be ~ 0.23 for neutron matter, if we use the calculations of Ref. [34].

Finally, we note that the calculation for the total susceptibility of Fantoni et al. [18] gives a result very close to the value one get by calculating G_0 from Brueckner theory and using Eq. (1) for the susceptibility, which does not take into account the renormalization of the magnetic moment. This would appear to indicate that the tensor correlations redistribute spectral weight between single pair and multipair excitations, leaving the total susceptibility unchanged. A similar effect is familiar for the compressibility of electrons in metals when one takes into account the electron-phonon interaction [24].

VI. SYMMETRIC NUCLEAR MATTER

Symmetric nuclear matter can be treated by a straightforward generalization of the discussion above by introducing the isospin degree of freedom in addition to the spin one. Thus the generalization of Eq. (5) becomes

$$\begin{aligned} f_{\mathbf{pp}'\sigma\sigma'} &= f_{\mathbf{pp}'} + f'_{\mathbf{pp}'}\boldsymbol{\tau} \cdot \boldsymbol{\tau}' + g_{\mathbf{pp}'}\boldsymbol{\sigma} \cdot \boldsymbol{\sigma}' + g'_{\mathbf{pp}'}\boldsymbol{\sigma} \cdot \boldsymbol{\sigma}'\boldsymbol{\tau} \cdot \boldsymbol{\tau}' \\ &+ f_{\mathbf{pp}'\sigma\sigma'}^T + f_{\mathbf{pp}'\sigma\sigma'}^{T'}\boldsymbol{\tau} \cdot \boldsymbol{\tau}', \end{aligned} \quad (79)$$

where $\boldsymbol{\tau}$ is the isospin operator. Since the magnetic moments of neutrons and protons are different, a magnetic field couples to both isoscalar and isovector components. Because the difference between the magnetic moments of a neutron and a proton is much larger than

TABLE III: Landau parameters for symmetric nuclear matter calculated by [5]. The Landau parameters are calculated for Fermi wave vector $k_F = 1.35 \text{ F}^{-1}$ and the effective mass is taken to be $m^*/m = 1$.

l	G_l	G'_l	$\tilde{H}_l$	$\tilde{H}'_l$
0	0.447	1.291	0.65	-0.255
1	0.760	0.070	0.975	-0.359
2	0.276	0.090	0.829	-0.315
3	0.078	0.069	0.617	-0.233

the sum, the magnetic response will be dominated by the isovector term. The isoscalar spin response is given by Eq. (22), but the density of states to be used is $N(0) = 2m^*p_F/\pi^2\hbar^3$, the factor of two reflecting the two nucleon species. The isovector spin response is given by a similar expression, but with G, H, K and L replaced by G', H', K' and L' . Since there is no calculation of the new tensor terms for symmetric nuclear matter, we estimate the matrix elements by using the exchange tensor Landau parameters for symmetric nuclear matter calculated in Ref. [5], which are listed in Table III. This gives the contribution to the matrix elements to be

$$\begin{aligned} \langle 01|A'|21 \rangle &\approx 0.11 \quad \text{and} \\ \langle 21|A'|21 \rangle &\approx 1.08. \end{aligned} \quad (80)$$

The calculations of Ref. [21] give for the anisotropic contribution to the isovector magnetic moment the value $\mu_T \approx 0.08 \mu_N$ and therefore the third and fourth terms in the analogue of Eq. (75) for the isovector magnetic response may be neglected. The second term is also unimportant, since it is on the order of 1% of the first term. We now use sum-rule arguments [19] to put a bound on the fifth term, which comes from multipair excitations. Equation (78) applies to a Fermi system with an arbitrary number of components, and recent calculations by Cowell and Pandharipande [34] of the static structure function and mean excitation energy for the isovector spin response. Their calculations give $S(q) = 0.155$ and $\bar{\omega} \approx 253$ MeV in the limit $q \rightarrow 0$. Comparing the contributions from multipair excitations with the susceptibility of a free Fermi gas we find

$$\frac{\chi_M(q, 0)}{\chi_F} \geq \frac{4\epsilon_F S(q)}{3\bar{\omega}}, \quad (81)$$

where $\chi_F = 3n/2\epsilon_F$ is the susceptibility of a free Fermi gas consisting of 2 species of spin-1/2 particles and ϵ_F is the Fermi energy. At nuclear matter density $n = 0.16 \text{ fm}^{-3}$, $\epsilon_F \approx 37$ MeV, gives that χ_M is at least $\sim 3\%$ of χ_F , which is much smaller than the corresponding bound for pure neutrons.

We are not aware of recent calculations of the total susceptibility for symmetric nuclear matter similar to the calculations of Ref. [18], but we note that for neutron matter those calculations gave a susceptibility very close

to the result one obtains by calculating G_0 from Brueckner theory and using Eq. (1) to calculate the susceptibility. Therefore we compare χ_M with the analogue for symmetric nuclear matter of Eq. (1), which we call χ_0 , and find

$$\frac{\chi_M}{\chi_0} \geq \frac{4\epsilon_F S(q)}{3\bar{\omega}} \frac{(1 + G'_0)}{m^*/m} \approx 0.09, \quad (82)$$

where we have taken $m^*/m = 0.8$, and G'_0 from Ref. [5], as listed in Table III. This result indicates that the multipair contributions, being at least 9 % of the total susceptibility, can have some importance for symmetric nuclear matter, even though the bound is much smaller than the corresponding bound for neutron matter. It would be valuable to have explicit calculations of the multipair contributions to the susceptibility since it is unclear to what extent the bound provides a realistic estimate of the value of the quantity.

VII. CONCLUDING REMARKS

In this paper we have derived an expression for the magnetic susceptibility of a Fermi liquid for a general interaction which conserves the total angular momentum and the total spin of the quasiparticle-quasihole pair. Apart from the contribution from single quasiparticle-quasihole pairs, which may be calculated using Landau theory, the susceptibility also contains a contributions from excitation of multipair states. In addition to the exchange tensor term usually included, we have taken into account the new tensor terms found in Ref. [7]. We have also introduced an alternative parametrization of the exchange tensor interaction which has the advantage of reducing the importance of high angular harmonic contributions.

For neutron matter we find that the most important effects of the tensor force are to renormalize the magnetic moment and to give a contribution from multipair states. For symmetric nuclear matter one can draw the same conclusion, the bound on the multipair contribution is, however, much smaller, but it is not clear how close the actual contribution to the susceptibility, from multipairs, is to this lower bound.

The formalism may easily be extended to calculate the response to weak probes: the only difference being that the magnetic moments must be replaced by the corresponding matrix elements of the weak charges.

Acknowledgments

We thank Achim Schwenk for interesting discussions and for providing us with his calculations of the tensor Landau parameters and Shannon Cowell and Vijay Pandharipande for giving us the result of their calculations of the static structure factor and energy-weighted sum. We are also grateful to Bengt Friman and Andrew Jackson for useful discussions and to Dan-Olof Riska for helpful correspondence. In addition, we thank ECT* and its director Wolfram Weise for hospitality in Trento. One of the authors (PH) was partially supported by the KBN grant no. 2P03D.020.20, and of us (EO) acknowledges financial support in part from a European Commission Marie Curie Training Site Fellowship under Contract No. HPMT-2000-00100. This work has been conducted within the framework of the school on Advanced Instrumentation and Measurements (AIM) at Uppsala University supported financially by the Foundation for Strategic Research (SSF).

-
- [1] A. B. Migdal, *Nuclear Theory: The Quasiparticle Method* (Benjamin, N. Y., 1968).
 - [2] J. Dąbrowski and P. Haensel, *Ann. Phys.* **97**, 452 (1976).
 - [3] J. Dąbrowski and P. Haensel, *Can. J. Phys.* **52**, 1768 (1974).
 - [4] P. Haensel and J. Dąbrowski, *Nucl. Phys. A* **254**, 211 (1975).
 - [5] S.-O. Bäckman, O. Sjöberg, and A. D. Jackson, *Nucl. Phys. A* **321**, 10 (1979).
 - [6] A. Schwenk, G. E. Brown, and B. Friman, *Nucl. Phys. A* **703**, 745 (2002).
 - [7] A. Schwenk and B. Friman, *Phys. Rev. Lett.* **92**, 082501 (2004).
 - [8] W. H. Dickhoff, A. Faessler, H. Münther, and S.-S. Wu, *Nucl. Phys. A* **405**, 534 (1983).
 - [9] H.-T. Janka, *Astrophys. J.* **368**, 527 (2001).
 - [10] R. F. Sawyer, *Phys. Rev. D* **11**, 2740 (1975).
 - [11] N. Iwamoto and C. J. Pethick, *Phys. Rev. D* **25**, 313 (1982).
 - [12] G. Raffelt and D. Seckel, *Phys. Rev. D* **52**, 1780 (1995).
 - [13] G. Raffelt, D. Seckel, and G. Sigl, *Phys. Rev. D* **54**, 2784 (1996).
 - [14] G. Raffelt and G. Sigl, *Phys. Rev. D* **60**, 023001 (1999).
 - [15] A. Burrows and R. F. Sawyer, *Phys. Rev. C* **58**, 554 (1998).
 - [16] M. Prakash, J. M. Lattimer, R. F. Sawyer, and R. R. Volkas, *Annu. Rev. Nucl. Part. Sci.* **51**, 295 (2001).
 - [17] S. Reddy, M. Prakash, and J. M. Lattimer, *Phys. Rev. D* **58**, 013009 (1998).
 - [18] S. Fantoni, A. Sarsa, and K. E. Schmidt, *Phys. Rev. Lett.* **87**, 181101 (2001).
 - [19] E. Olsson and C. J. Pethick, *Phys. Rev. C* **66**, 065803 (2002).
 - [20] H. Miyazawa, *Prog. Theor. Phys.* **6**, 801 (1951).
 - [21] A. Arima, K. Shimizu, W. Bentz, and H. Hyuga, *Adv. Nucl. Phys.* **18**, 1 (1987).
 - [22] K. Tsushima, D. O. Riska, and P. G. Blunden, *Nucl. Phys. A* **559**, 543 (1993).

- [23] P. Haensel and A. J. Jerzak, Phys. Lett. B **112**, 285 (1982).
- [24] A. J. Leggett, Phys. Rev. **140**, 1869 (1965).
- [25] G. Baym and C. J. Pethick, *Landau Fermi-liquid theory: concepts and applications* (Wiley N. Y., 1991), p. 71.
- [26] L. D. Landau, Sov. Phys. JETP **3**, 920 (1957).
- [27] L. D. Landau, Sov. Phys. JETP **5**, 101 (1957).
- [28] C. J. Pethick and G. M. Carneiro, Phys. Rev. A **7**, 304 (1973).
- [29] C. J. Pethick and G. M. Carneiro, Phys. Rev. B **11**, 1106 (1975).
- [30] A. Schwenk (2003), private communication.
- [31] A. Schwenk, B. Friman, and G. E. Brown, Nucl. Phys. A **713**, 191 (2003).
- [32] S. Cowell and V. R. Pandharipande, Phys. Rev. C **67**, 035504 (2003).
- [33] A. Akmal and V. R. Pandharipande, Phys. Rev. C **56**, 2261 (1997).
- [34] S. Cowell and V. R. Pandharipande (2004), private communication.